

FIG. 1




FIG. 2






FIG. 3



**Key:** a) n-BuLi, Ether; Et<sub>2</sub>AlCl, toluene; b) 5 then dil.HCl; c) Lindlar Catalyst, H<sub>2</sub>; d) TsCl, THF, pyridine; e) TMStriflate, CH<sub>2</sub>Cl<sub>2</sub>, 2,6-Lutidine; f) NaI, acetone, Δ; g) N-Propionylcamphorsultam 18, n-BuLi, then 17; h) DIBAH, THF-CH<sub>2</sub>Cl<sub>2</sub>.

FIG. 4



**Key:** a) MEMCl, DIPA, CH<sub>2</sub>Cl<sub>2</sub>; b) TBAF, THF; c) Swern Oxidation; d) Horner-Emmons Reaction, LDA, THF, 24; then ketone 23; e) HCl, H<sub>2</sub>O, THF; f) TsCl, pyridine, CH<sub>2</sub>Cl<sub>2</sub>; g) TBSOTf, DIPA; h) NaI, acetone, Δ; i) N-propionylcamphorsultam 18, n-BuLi, then iodide; j) DIBAH, CH<sub>2</sub>Cl<sub>2</sub>.

FIG. 5




FIG. 5a




FIG. 6





FIG. 7



Key: a)  $\text{Cl}_3\text{C}_6\text{H}_2\text{COCl}$ , pyridine, DMAP; b) TBAF, THF; c) PCC,  $\text{CH}_2\text{Cl}_2$ ; d) Horner-Emmons: LDA, 24.

FIG. 8





FIG. 9



Reported, Nicolaou, +16.1  
Found, -17.6

FIG. 10








FIG. 11




Key: a) **58**, TBAF, THF; **58a**, chromatography; **58b**, dil. acid or DDQ, CH<sub>2</sub>Cl<sub>2</sub>, water; b) PhSO<sub>2</sub>Cl, pyridine, or Cl<sub>3</sub>PhCOCl, pyridine, DMAP, CH<sub>2</sub>Cl<sub>2</sub>.



Key: a) as in Figure 7




**Key:** a) 1.0 TBAF, THF; b) PCC,  $\text{CH}_2\text{Cl}_2$ ; c) pyridine or DMAP,  $\text{CH}_2\text{Cl}_2$ ; d) Horner-Emmons: LDA, **24** or other phosphonates.




**Key:** a) i) Mg, ether; ii)  $\text{CuBr}\cdot\text{DMS}$ ; iii) propyne; iv)  $\text{I}_2$ ; b) i)  $n\text{-BuLi}$ ; ii)  $\text{Me}_2\text{AlCl}$ ; iii) **5b**; c) HCl, EtOH; d) TsCl, pyridine; e) TBSOTf, 2,6-lutidine,  $\text{CH}_2\text{Cl}_2$ ; f) NaI, acetone; g) **18**,  $n\text{-BuLi}$ ,  $-40^\circ\text{C}$ , THF; h)  $\text{LiAlH}_4$ , THF; i) pyridine- $\text{SO}_3$ ,  $\text{CH}_2\text{Cl}_2$ ;  $\text{Et}_3\text{N}$ .





**Key:** a) i) Mg, ether; ii) CuBr·DMS, DMS, ether; iii) propyne; iv) pentynyl lithium; b) 5b, -40 °C, 36 hrs; c) TBSOTf, 2,6-lutidine, CH<sub>2</sub>Cl<sub>2</sub>; d) AD-mix a, e) NaIO<sub>4</sub>, EtOH, HOH; f) NaBH<sub>4</sub>, MeOH; g) TsCl, pyridine.



**Key:** a) NaI, acetone; b) 18, n-BuLi, -40 °C, THF; c) LiAlH<sub>4</sub>, THF; d) pyridine·SO<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>; Et<sub>3</sub>N; e) DIBAL, ether, -78 °C.






FIG. 19




Key: a) PCC,  $\text{CH}_2\text{Cl}_2$ ; b) pyridine or DMAP,  $\text{CH}_2\text{Cl}_2$ ; c) Horner-Emmons: LDA, **24** or other phosphonates.

FIG. 20



Key: a)  $\text{O}_3$ , MeOH,  $-78^\circ\text{C}$ ; then  $\text{NaBH}_4$ ; b)  $\text{TsCl}$ , pyridine; c)  $\text{LiBr}$ , acetone; d) i)  $\text{Mg}$ , ether; ii)  $\text{CuBr}\cdot\text{DMS}$ , DMS, ether; iii) propyne; iv) pentynyl lithium; v) **5a** or **5b**,  $-40^\circ\text{C}$ , 36 hrs; e)  $\text{TMSOTf}$ , 2,6-lutidine,  $\text{CH}_2\text{Cl}_2$ ; or  $p\text{-MeOC}_6\text{H}_4\text{CH}_2\text{Br}$ , NaH, DMF; f) ADMix a; then  $\text{NaIO}_4$ .

FIG. 21



**Key:** a) i) Mg, ether; ii) CuBr•DMS, DMS, ether; iii) propyne; iv) pentynyl lithium; v) **5a**, -40 °C, 36 hrs; b) i) Mg, ether; ii) CuBr•DMS, DMS, ether; iii) propyne; iv) pentynyl lithium; v) **5a**, -40 °C, 36 hrs; vi) TMSOTf, -78 °C; c) TMSOTf, 2,6-lutidine, CH<sub>2</sub>Cl<sub>2</sub>; d) (ipc)<sub>2</sub>BH, THF, -20 °C; then H<sub>2</sub>O<sub>2</sub>, NaOH; e) pyr•SO<sub>3</sub>, DMSO, Et<sub>3</sub>N, CH<sub>2</sub>Cl<sub>2</sub>; f) (ipc)<sub>2</sub>BH, THF, -20 °C; then PCC, CH<sub>2</sub>Cl<sub>2</sub>.

FIG. 22






FIG. 23




Key: a) Cr(VI), or pyridine $\cdot$ SO<sub>3</sub>, DMSO, Et<sub>3</sub>N, CH<sub>2</sub>Cl<sub>2</sub>; b) LDA, 24, then 140; c] i) 123, Mg, ether; ii) CuBr $\cdot$ DMS, DMS, ether; iii) propyne; iv) pentynyl lithium; v) 141, -40 °C, 36 hrs; vi) TMSOTf, -78 °C; d) (ipc)<sub>2</sub>BH; then Cr(VI); e) 56a, THF, -78 °C; then silica gel; f) PhSO<sub>2</sub>Cl, pyridine, CH<sub>2</sub>Cl<sub>2</sub>; g) dimethyldioxirane, acetone; h) chiral ketone 145, oxone, pH 7-8, aq. CH<sub>3</sub>CN (Y. Shi, et al., J. Org. Chem., 63(23), 8475 (1998)).

FIG. 24



Key: a) AD-mix; then NaIO<sub>4</sub>; b) NaBH<sub>4</sub>, MeOH; c) TBSCl, pyridine, CH<sub>2</sub>Cl<sub>2</sub>; d) O<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>; Me<sub>2</sub>S; e) Ph<sub>3</sub>P=CH-I, THF; f) *t*-BuLi, then Et<sub>2</sub>AlCl, then 141, then TIPSCl; g) Quinolinium fluorochromate, CH<sub>2</sub>Cl<sub>2</sub>; h) 56a, THF, -78 °C; i) HF $\cdot$ pyr, CH<sub>3</sub>CN; j) PhSO<sub>2</sub>Cl, pyridine, CH<sub>2</sub>Cl<sub>2</sub>; k) dimethyldioxirane, acetone.



Key: a) Zn/Cu, sonochem; b) Ph<sub>3</sub>P=CH-I, THF; c) (Ipc)<sub>2</sub>BH; then NaBO<sub>3</sub>; d) TBSCl, pyr, CH<sub>2</sub>Cl<sub>2</sub>.






FIG. 26b



**Key:** a) Mg, ether; then propyne, Cu(I); pentynyl lithium; epoxide **143** (Scheme XXIII); then  $\text{H}_3\text{O}^+$ ; b)  $\beta$ -lactone **157** (Scheme XXIII), pyridine,  $\text{CH}_2\text{Cl}_2$ ; c) Mg, ether; then propyne, Cu(I); pentynyl lithium;  $\beta$ -lactone **157** (Scheme XXIII); then  $p\text{-MeOC}_6\text{H}_4\text{CH}_2\text{OCH}_2\text{Cl}$  or other protecting group such as TBSOTf or TBSCl; d)  $(\text{Ipc})_2\text{BH}$ , THF; then  $\text{LiOOH}$ ; e) Swern Oxidation; f)  $(\text{Ipc})_2\text{BH}$ , THF; then PCC; g) Lewis or protic acid; or alternatively base catalyzed cyclization; h) DDQ,  $\text{CH}_2\text{Cl}_2$ , HOH, buffer to remove the PMB, PMBM or DMB groups; Fluoride ion to remove Si based groups.




FIG. 27



**Key:** a) RCOX, pyridine, catalytic DMAP, CH<sub>2</sub>Cl<sub>2</sub>; b) TMSOTf, 2,6-lutidine, CH<sub>2</sub>Cl<sub>2</sub>; c) R<sub>1</sub>COX, DMAP, CH<sub>2</sub>Cl<sub>2</sub>; d) R<sub>1</sub>COX, DMAP, CH<sub>2</sub>Cl<sub>2</sub>; then silica gel. Where RCOX = active ester of usual variety.


[illegible]

FIG. 29




**Key:** a) i. Mg, ether; ii. CuBr•DMS complex, ether, DMS, -45°C; iii. propyne, -23°C; iv. Li-hexyne, HMPA, -78°C; v. 5c, -78°C to -25°C, 78%; b) SEM-Cl, DIPEA, CH<sub>2</sub>Cl<sub>2</sub>, 94%; c) DDQ, CH<sub>2</sub>Cl<sub>2</sub>, HOH; 85%; d) SO<sub>3</sub>•pyr, TEA, CH<sub>2</sub>Cl<sub>2</sub>; 78%; e) 24, n-BuLi, THF, -78°C to R.T., 85%; f) (Ipc)<sub>2</sub>BH, THF; H<sub>2</sub>O<sub>2</sub>, 82%; g) oxalyl chloride, DMSO, TEA, CH<sub>2</sub>Cl<sub>2</sub>, 88%; h) LDA, -78°C to -40°C, ZnCl<sub>2</sub>; -78°C to -50°C, THF, 68%; i) TROC-Cl, DMAP, CH<sub>2</sub>Cl<sub>2</sub>, j) TFA, CH<sub>2</sub>Cl<sub>2</sub>, k) Trichlorobenzoyl chloride, TEA, THF, DMAP, toluene; l) HF•pyridine; m) Zn, HOAc; n) m-CPBA, CH<sub>2</sub>Cl<sub>2</sub>.

FIG. 30



Key: a) HF•pyridine, THF; b) Zn, HOAc; c) RCOOH, DCC, TEA, DCM.

FIG. 31



**Key:** a) LDA,  $-78^{\circ}\text{C}$  to  $-40^{\circ}\text{C}$ ,  $\text{ZnCl}_2$ ;  $-78^{\circ}\text{C}$  to  $-50^{\circ}\text{C}$ , THF, 68%; then  $\text{CH}_2=\text{CH}(\text{CH}_2)_3\text{COCl}$ , DMAP,  $\text{CH}_2\text{Cl}_2$ , b) TFA,  $\text{CH}_2\text{Cl}_2$ , c) Trichlorobenzoyl chloride, TEA, THF, DMAP, toluene; d)  $\text{HF}\cdot\text{pyridine}$ ; e) vicinal dihydroxylation; f)  $\text{NaIO}_4$ , THF, HOH.

FIG. 32




FIG. 33






FIG. 34




Key: a) PMB-Br, NaH, Bu<sub>4</sub>N-I, THF, 0 °C, 85%; b) i) Mg, ether, rt; ii) CuBr-DMS, ether, DMS, -45 °C, 3h, iii) Propyne, -45 °C to -23 °C, 4h then lithiohexyne, -78 °C, 1h; iv) epoxide 205, -78 °C, 1h, -25 °C, 24h, 76%; c) SEMCl, DIPEA, DCM, 0 °C, 92%; d) DDQ, DCM:water (8:2), 88%; e) DMSO, (COCl)<sub>2</sub>, DCM, TEA, -78 °C, 85%; f) 207, *n*-BuLi, THF, then 214, 72%; g) (*i*-PC)<sub>2</sub>BH, THF, 0.5h, aq. NaBO<sub>3</sub>; and h) DMSO, (COCl)<sub>2</sub>, DCM, TEA, -78 °C, 92%.

FIG. 35




**Key:** (a) (i)  $\text{Bu}_2\text{BOTf}$ , DIPEA,  $\text{CH}_2\text{Cl}_2$ ,  $0^\circ\text{C}$  then add **217** at  $-78^\circ\text{C}$ ; (ii) Raney Ni, acetone,  $60^\circ\text{C}$ , 45 min, 70% combined; (b) (i) TBDMSOTf, 2,6-lutidine,  $\text{CH}_2\text{Cl}_2$ ,  $0^\circ\text{C}$  to rt, 95%; (ii) LiOH,  $\text{H}_2\text{O}_2$ , THF- $\text{H}_2\text{O}$ , rt, 82%.

FIG. 36



**Key:** a) LDA, **204**, THF,  $-78^\circ\text{C}$  to  $-40^\circ\text{C}$  then to  $-78^\circ\text{C}$ ,  $\text{ZnCl}_2$ , **203**,  $-78^\circ\text{C}$  to  $-50^\circ\text{C}$ , 0.5h; b) TrocCl, Py, DCM,  $0^\circ\text{C}$ ; c) TFA, DCM (3:7),  $-20^\circ\text{C}$ , 1h, 63% (three steps).





**Key:** (a) 2,4,6-Cl<sub>3</sub>C<sub>6</sub>H<sub>2</sub>COCl, TEA, THF, DMAP, toluene, rt, 1h; b) HF-Py, DCM, rt, 95%;  
c) Zn, aq. NH<sub>4</sub>Cl, MeOH, reflux, 92%; d) [Methyl(trifluoromethyl)]dioxirane, MeCN, 0 °C, 56%.